arXiv:1304.2519v1 [cond-mat.stat-mech] 9 Apr 2013

Nonlinear subdiffusive fractional equations and aggregation phenomenon

Sergei Fedotov
School of Mathematics, The University of Manchester, Manchester M13 9PL, UK

In this article we address the problem of the nonlinear interaction of subdiffusive particles. We introduce the random walk model in which statistical characteristics of a random walker such as escape rate and jump distribution depend on the mean field density of particles. We derive a set of nonlinear subdiffusive fractional master equations and consider their diffusion approximations. We show that these equations describe the transition from an intermediate subdiffusive regime to asymptotically normal advection-diffusion transport regime. This transition is governed by nonlinear tempering parameter that generalizes the standard linear tempering. We illustrate the general results through the use of the examples from cell and population biology. We find that a nonuniform anomalous exponent has a strong influence on the aggregation phenomenon.

PACS numbers:

I. INTRODUCTION

Anomalous subdiffusion is a widespread phenomenon in physics and biology [1–3]. It is observed in the transport of proteins and lipids on cell membranes [4], RNA molecules in the cells [5], signaling molecules in spiny dendrites [6], and elsewhere. Apart from fractional Brownian motion, the *linear* fractional equations are the standard models for the description of anomalous subdiffusive transport [2]. In these models the diffusing particles do not interact. The question then arises as to how to extend these equations for the *nonlinear* case, involving particles interactions. These nonlinear effects are typical and very important in cellular and population biology. It is well known that many biophysical processes in microorganisms depend on the their population density. A typical example is the *quorum sensing phenomenon,* in which microorganisms coordinate their behaviour according to their local population density [7]. Other examples are a *cellular adhesion,* which involves the interaction between neighbouring cells [8–11]; and the *volume-filling effect,* which describes the dependence of cell motility on the availability of space in a crowded environment [12]. The understanding of macroscopic phenomena like cell and microorganizm aggregation requires an understanding of how individual species interact through attractive or repulsive forces (see, for example, [13] and references therein). The attraction between individuals may result from various social interactions such as mating, settlement, defense against predators, etc. While the repulsion may occur due to low resources in highly populated regions [13, 14]. Note that microorganisms interact both directly and indirectly via signaling molecules.

The main purpose of this paper is to incorporate these nonlinear effects into subdiffusive equations. Our aim is to take into account the interaction between particles on the mesoscopic level, at which the random walker's characteristics depend on the mean field density of particles. Our intention is to derive the nonlinear subdiffusive fractional equations for the density of particles and apply these equations to the problem of aggregation. In this paper we use two different approaches that are based on the density-dependent dispersal kernels and density-dependent jump rate. Note that several theoretical studies have been devoted to nonlinear generalizations of linear subdiffusive fractional equations. However, most research has been focused on the problem how to incorporate the nonlinear reactions into subdiffusive equations [15–24]. The aim of this paper is to study the nonlinear subdiffusive transport processes involving the anomalous trapping of particles and their interactions.

In this paper we deal with the random walk model involving a residence time dependent escape rate and the structural density of particles. This has been used by many authors for the analysis of the non-Markovian random walks [3, 15, 16, 25, 26]. It turns out that this linear model is the most suitable for further nonlinear generalizations. We consider a 'space-jump' random walk in one space dimension. The particle waits for a random time (residence time) T_x at point x in space before making a jump to another point. The random residence time T_x is determined by the probability density function $\psi(x,\tau) = \Pr\{\tau < T_x < \tau + d\tau\}$. The key characteristic of this random walk is the escape rate γ from the point x. It depends on the residence time τ and the position $x : \gamma = \gamma(x,\tau)$. This rate can be rewritten in terms of the probability density function $\psi(x,\tau)$ and the survival probability $\Psi(x,\tau) = \int_t^\infty \psi(x,u)du$ as follows [27]

$$\gamma(x,\tau) = \frac{\psi(x,\tau)}{\Psi(x,\tau)}. \tag{1}$$

Let $\xi(x,\tau,t)$ be the structural density of particles at point x at time t whose residence time lies in the interval $(\tau,\tau+d\tau)$. We assume that at $t=0$ all particles have zero residence time: $\xi(x,\tau,0) = \rho_0(x)\delta(\tau)$. The mean escape rate, $i\,(x,t)$, of particles from the point x can be defined as

$$i(x,t) = \int_0^t \gamma(x,\tau)\xi(x,\tau,t)d\tau \tag{2}$$

and the density of particles $\rho\,(x,t)$

$$\rho\,(x,t) = \int_0^t \xi(x,\tau,t)d\tau. \tag{3}$$

One of the main results in the non-Markovian random walk theory is that the mean escape rate $i\left(x,t\right)$ can be written as a convolution [3]

$$i\left(x,t\right)=\int_0^t K\left(x,t-\tau\right)\rho\left(x,\tau\right)d\tau. \tag{4}$$

Here $K(x,t)$ is the memory kernel defined by its Laplace transform

$$\hat{K}\left(x,s\right)=\frac{\hat{\psi}\left(x,s\right)}{\hat{\Psi}\left(x,s\right)}, \tag{5}$$

where $\hat{\psi}\left(x,s\right) = \int_0^\infty e^{-st}\psi(x,\tau)d\tau$ and $\hat{\Psi}\left(x,s\right) = \int_0^\infty e^{-st}\Psi(x,\tau)d\tau$ [3]. The master equation for the density $\rho\left(x,t\right)$ can be written as the balance of particles at the point x

$$\frac{\partial\rho}{\partial t}=\int_{\mathbb{R}} i\left(x-z,t\right)w\left(z|x-z\right)dz-i\left(x,t\right), \tag{6}$$

where $w\left(z|x\right)$ is the dispersal kernel.

In the anomalous subdiffusive case, the survival probability $\Psi_\mu\left(x,t\right)$ can be modelled by the Mittag-Leffler function [28]

$$\Psi_\mu\left(x,t\right)=E_\mu\left[-\left(\frac{\tau}{\tau_0(x)}\right)^{\mu(x)}\right],\ 0<\mu(x)<1, \tag{7}$$

where $\mu(x)$ is the space-dependent anomalous exponent. In what follows we assume for simplicity that τ_0 is constant. The Laplace transform of the memory kernel $K_\mu\left(x,t\right)$ is $\hat{K}_\mu\left(x,s\right) = s^{1-\mu(x)}{\tau_0}^{-\mu(x)}$ and the integral anomalous escape rate $i\left(x,t\right)$ can be written as

$$i\left(x,t\right)=\frac{1}{{\tau_0}^{\mu(x)}}\mathcal{D}_t^{1-\mu(x)}\rho\left(x,t\right), \tag{8}$$

where $\mathcal{D}_t^{1-\mu(x)}$ is the The Riemann-Liouville derivative with varying anomalous exponent [29]. Substitution of (8) into (6) gives the integral fractional equation with space-dependent anomalous exponent

$$\begin{aligned}\frac{\partial\rho}{\partial t} &= \int_{\mathbb{R}}\frac{1}{{\tau_0}^{\mu(x-z)}}\mathcal{D}_t^{1-\mu(x-z)}\rho\left(x-z,t\right)w\left(z|x-z\right)dz\\ &-\frac{1}{{\tau_0}^{\mu(x)}}\mathcal{D}_t^{1-\mu(x)}\rho\left(x,t\right).\end{aligned} \tag{9}$$

Using the Taylor series expansion in terms of z, and a symmetric dispersal kernel w, we obtain the subdiffusive fractional equation

$$\frac{\partial\rho}{\partial t}=\frac{\partial^2}{\partial x^2}\left[D_\mu\left(x\right)\mathcal{D}_t^{1-\mu(x)}\rho\left(x,t\right)\right], \tag{10}$$

with the fractional diffusion coefficient $D_\mu(x) = \sigma^2(x)/{\tau_0}^{\mu(x)}$, where $\sigma^2(x)=\int z^2 w(z|x)dz$.

It has been found recently that subdiffusive fractional equations are not structurally stable with respect to the spatial variations of fractal exponent $\mu(x)$. This leads to the anomalous aggregation of particles [26]. In heterogeneous biological systems, in which the exponent $\mu(x)$ is space-dependent, the question arises as to whether this anomalous aggregation of a population can be prevented. Therefore it is an important problem to find the way how to regularize subdiffusive fractional equations. One way is to incorporate random killing, which ensures regular behaviour in the long-time limit [30]. The aim of this paper is to address this problem through a nonlinear escape rate that takes into account repulsive forces between particles. In the next section we derive the nonlinear generalization of the fractional equations like (9) and (10).

II. NONLINEAR SUBDIFFUSIVE FRACTIONAL EQUATIONS

There are two major ways in which nonlinear density-dependence effects can be implemented into non-Markovian and subdiffusive transport equations. The simplest way is to take into account the dependence of jump density w on ρ. This dependence can take into account various non-linear effects such as adhesion, quorum sensing, volume filling, etc. However, we begin with more complicated case of the random walk model for which the escape rate is a function of the residence time and the local density of particles.

A. Nonlinear escape rate with memory and nonlinear tempering

The main problem with this anomalous escape rate (8) is the phenomenon of anomalous aggregation [25, 26]. Nonuniform distribution of the anomalous exponent $\mu(x)$ over the finite domain $[0,L]$ leads to

$$\rho\left(x,t\right)\to\delta(x-x_M)\qquad as\qquad t\to\infty. \tag{11}$$

Here x_M is the point in space where the anomalous exponent $\mu\left(x\right)$ has a minimum. The problem is that that the escape rate $i\left(x,t\right)$ is a linear functional of the density of particles $\rho\left(x,t\right)$ and does not take into account nonlinear effects of repulsive forces which, in many situations, can prevent anomalous aggregation. According to (11) all particles aggregate into a small region around the point $x=x_M$ forming a high density system. To prevent such anomalous aggregation, one can assume that the overcrowding leads to an increase of repulsive forces and a corresponding correction of the anomalous escape rate $\gamma(x,\tau)$.

We assume that the probability of escape due to the repulsive forces is independent from anomalous trapping. We define this probability for a small time interval Δt as

$$\alpha(\rho(x,t))\Delta t+o(\Delta t). \tag{12}$$

Here $\alpha(\rho)$ is the transition rate which is an increasing function of the particles density ρ. Another interpretation can be given in terms of the quorum sensing phenomenon [7]. The large cell density can lead to the local over-depletion of nutrients and oxygen and as a result cells can change phenotype from a proliferating state to a migrating one [21]. In another words, when the concentration of cells is low, $\alpha(\rho) = 0,$ but if the concentration of cells $\rho(x,t)$ reaches a certain level $\rho_{cr} : \alpha(\rho) \neq 0$ for $\rho \geq \rho_{cr}$. Of course, one can assume a non-monotonic dependence of the transition rate $\alpha(\rho)$ on ρ. For example, at low cell densities $\alpha(\rho)$ could decrease with ρ, while at high densities $\alpha(\rho)$ could be an increasing function: $\alpha(\rho) = \alpha_0 \left(1 - a_1\rho + a_2\rho^2\right)$ [12, 13].

Firstly we formulate the Markovian model for the random walk of particles with the density-dependent escape rate. The balance equation for the structural density $\xi(x,\tau,t)$ takes the form

$$\begin{aligned}\xi(x,\tau+\Delta t,t+\Delta t,) &= \xi(x,\tau,t)\left(1-\gamma(x,\tau)\Delta t\right)\times\\ &\quad\left(1-\alpha(\rho(x,t))\,\Delta t\right)+o\left(\Delta t\right).\end{aligned}$$

where $1 - \gamma(x,\tau)\Delta t$ is the survival probability during time Δt due to the trapping, and $1-\alpha(\rho(x,t))\Delta t$ is the survival probability during time Δt corresponding to the repulsion forces between particles. In the limit $\Delta t \to 0$, we obtain

$$\frac{\partial \xi}{\partial t} + \frac{\partial \xi}{\partial \tau} = -\left[\gamma(x,\tau) + \alpha(\rho(x,t))\right]\xi, \tag{13}$$

where the effective transition rate is the sum of two escape rates:

$$\gamma(x,\tau) + \alpha(\rho(x,t)). \tag{14}$$

The second term $\alpha(\rho(x,t))$ can be treated as the correction of the basic escape rate $\gamma(x,\tau)$ due to repulsive forces. The boundary condition for $\xi(x,\tau,t)$ at zero residence time $\tau = 0$ is

$$\begin{aligned}\xi(x,0,t) &= \int_{\mathbb{R}}\int_0^t \left[\gamma(x-z,\tau)+\alpha(\rho(x-z,t))\right]\times\\ &\quad\xi(x-z,\tau,t)w\left(z|\rho(x-z,t)\right)d\tau dz. \qquad (15)\end{aligned}$$

To obtain the effective escape rate

$$i\left(x,t\right) = \int_0^t \gamma(x,\tau)\xi(x,\tau,t)d\tau + \alpha(\rho)\rho(x,t) \tag{16}$$

we solve (13) by the methods of characteristics and substitute $\xi(x,\tau,t)$ into (15) and (16). Finally we obtain

$$\begin{aligned}i\left(x,t\right) &= \int_0^t K\left(x,t-\tau\right)e^{-\int_\tau^t \alpha(\rho(x,s))ds}\rho\left(x,\tau\right)d\tau\\ &\quad+\alpha(\rho)\rho(x,t). \qquad (17)\end{aligned}$$

One can see that the first term in (17) involves the exponential factor, $e^{-\int_\tau^t \alpha(\rho(x,s))ds}$, which can be interpreted as a nonlinear tempering. The main feature of this modified escape rates is that although the *local* rates $\gamma(x,\tau)$ and $\alpha(\rho(x,t))$ are additive (see (14)), the corresponding terms in the *integral* escape rate (17) are not additive. This is clearly non-Markovian effect. One of the main aims of this paper is to find out what the implications of this effect on the long-time behaviour of the density $\rho(x,t)$ are (see section III).

Note that when γ does not depend on the residence time variable τ, the survival function has an exponential form $\Psi(x,\tau) = e^{-\gamma(x)\tau}$, the Laplace transform $\hat{K}\left(x,s\right) = \gamma(x)$ and the Markovian escape rate

$$i(x,t) = \left[\gamma(x) + \alpha(\rho)\right]\rho\left(x,t\right). \tag{18}$$

Substitution of this into (6) gives the closed nonlinear Kolmogorov-Feller equation for $\rho\left(x,t\right)$ [3]. One can find various nonlinear diffusion approximations of this equation assuming density-dependent dispersal kernels $w\left(z|\rho\left(x,t\right)\right)$ and density-dependent jump rate $\alpha(\rho\left(x,t\right))$ [13].

B. Anomalous subdiffusive case

In the subdiffusive case, it follows from (17) that instead of (8) we have

$$\begin{aligned}i\left(x,t\right) &= \frac{e^{-\Phi(x,t)}}{{\tau_0}^{\mu(x)}}\mathcal{D}_t^{1-\mu(x)}\left[e^{\Phi(x,t)}\rho\left(x,t\right)\right]\\ &\quad+\alpha(\rho)\rho(x,t), \qquad (19)\end{aligned}$$

where

$$\Phi\left(x,t\right) = \int_0^t \alpha\left(\rho\left(x,s\right)\right)ds. \tag{20}$$

This function plays a very important role in what follows. For the jump density involving only two outcomes

$$w\left(z|x,t\right) = r(x,t)\delta(z-a) + l(x,t)\delta(z+a), \tag{21}$$

where a is the jump size, we obtain the master equation for $\rho\left(x,t\right)$ in terms of the total flux $i\left(x,t\right)$

$$\begin{aligned}\frac{\partial \rho}{\partial t} &= r(x-a,t)i(x-a,t)\\ &\quad+l(x+a,t)i(x+a,t) - i(x,t). \qquad (22)\end{aligned}$$

Here $r(x,t)$ is the jump probability from the point x to $x+a$, $l(x,t)$ is the jump probability from the point x to $x-a$. One can introduce the potential $U(x,t)$ that induce the bias of the random walk in one direction [31, 32]. In the case of chemotaxis, $U(x,t)$ can be interpreted as the density of chemotactic substance [33, 34]. In the limit $a \to 0$, by using

$$r(x-a,t) - l(x+a,t) = -\beta a\frac{\partial U}{\partial x} + o(a), \tag{23}$$

we obtain

$$\frac{\partial \rho}{\partial t} = -\beta a^2 \frac{\partial}{\partial x}\left[\frac{\partial U}{\partial x}i(x,t)\right] + a^2\frac{\partial^2 i(x,t)}{\partial x^2}. \tag{24}$$

This equation can be rewritten in the form of a nonlinear fractional Fokker-Planck equation

$$\frac{\partial \rho}{\partial t} = -\beta a^2 \frac{\partial}{\partial x}\left[\frac{\partial U}{\partial x}\left(\frac{e^{-\Phi}}{{\tau_0}^{\mu(x)}}\mathcal{D}_t^{1-\mu(x)}[e^{\Phi}\rho] + \alpha(\rho)\rho\right)\right] + a^2 \frac{\partial^2}{\partial x^2}\left[\frac{e^{-\Phi}}{2{\tau_0}^{\mu(x)}}\mathcal{D}_t^{1-\mu(x)}[e^{\Phi}\rho] + \alpha(\rho)\rho\right]. \qquad (25)$$

This equation is one of the main results of this paper. It describes the transition from subdiffusive transport to asymptotic normal advection-diffusion transport. We discuss this asymptotic transition in section III. Note that the above equation can be rewritten in terms of the fractional diffusion coefficient

$$D_\mu(x) = \frac{a^2}{2{\tau_0}^{\mu(x)}} \qquad (26)$$

which is finite in the limit $a \to 0$ and $\tau_0 \to 0$.

C. Nonlinear jump distributions

In this subsection we discuss various nonlinear jump distributions leading to nonlinear fractional equations. As we mentioned previously, this is the simplest way to incorporate nonlinearity into subdiffusive fractional equations [26]. As long as the escape rate $i\,(x,t)$ is determined, we can define "where to jump" through the dispersal kernel. Let us assume that the dispersal kernel w depends on the density of particles: $w = w(z|\rho(x,t))$. In what follows we restrict ourselves to the subdiffusive case for which the rate $i(x,t)$ is determined by (8).

1. *Local nonlinearity*

First we consider the case for which the jump dependence on ρ is local. For example, one can use Gaussian dispersal kernel with rapidly decaying tails as

$$w\,(z|\rho) = \frac{1}{\sqrt{2\pi\sigma^2(\rho)}}\exp\left[-\frac{z^2}{2\sigma^2(\rho)}\right].$$

An increasing dispersion $\sigma^2(\rho)$ describes the effect of the local repulsive forces, while a decaying $\sigma^2(\rho)$ corresponds to attractive forces. Using the Taylor series expansion in terms of z in the master equation (6) with $w\,(z|\rho(x-z,t))$, we obtain the subdiffusive equation

$$\frac{\partial \rho}{\partial t} = \frac{\partial^2}{\partial x^2}\left[K_\mu\,(\rho)\,\mathcal{D}_t^{1-\mu(x)}\rho\,(x,t)\right] \qquad (27)$$

with the nonlinear fractional diffusion coefficient

$$D_\mu\,(\rho) = \frac{\sigma^2(\rho)}{{\tau_0}^{\mu(x)}}.$$

One can also introduce the nonlinear drift term generated by Bernoulli jump distribution [13]

$$\begin{aligned} w\,(z|\rho) \; &= \; \frac{1}{2}\left(1 + au\,(\rho)\right)\delta\,(z-a) + \\ &\quad\; \frac{1}{2}\left(1 - au\,(\rho)\right)\delta\,(z+a)\,, \end{aligned}$$

where the positive function $u\,(\rho)$ takes into account the fact that the jump on the right is more likely than a jump to the left. It is assumed that $1 - au\,(\rho)$ is not negative; that is, $u\,(\rho) \leq a^{-1}$. The fractional master equation takes the form

$$\begin{aligned} \frac{\partial \rho}{\partial t} \; &= \; \frac{1 + au\,(\rho\,(x-a,t))}{2{\tau_0}^{\mu(x-a)}}\mathcal{D}_t^{1-\mu(x-a)}\rho\,(x-a,t) + \\ &\quad\; \frac{1 - au\,(\rho\,(x+a,t))}{2{\tau_0}^{\mu(x+a)}}\mathcal{D}_t^{1-\mu(x+a)}\rho\,(x+a,t) \\ &\quad\; -\frac{1}{{\tau_0}^{\mu(x)}}\mathcal{D}_t^{1-\mu(x)}\rho\,(x,t)\,. \end{aligned} \qquad (28)$$

Using the Taylor series expansion in (28), we obtain the fractional Fokker-Planck equation

$$\frac{\partial \rho}{\partial t} = -\frac{\partial}{\partial x}\left[2D_\mu(x)u\,(\rho\,(x,t))\,\mathcal{D}_t^{1-\mu(x)}\rho\right] + \frac{\partial^2}{\partial x^2}\left[D_\mu(x)\mathcal{D}_t^{1-\mu(x)}\rho\right] \qquad (29)$$

with the *nonlinear* advection term and fractional diffusion coefficient $D_\mu(x)$.

2. *Nonlocal nonlinearity*

Here we consider the volume filling effect and cell-to-cell adhesion [12]. One can use the jump density

$$w\,(z|\rho) = r(\rho)\delta\,(z-a) + l(\rho)\delta\,(z+a)\,. \qquad (30)$$

where $r(\rho)$ is the probability of jumping from x to $x+a$ and $l(\rho)$ is the probability of jumping from x to $x-a$ defined as follows

$$r(\rho) = \frac{\lambda(\rho)}{\lambda(\rho)+\mu(\rho)}, \quad l(x|\rho) = \frac{\mu(\rho)}{\lambda(\rho)+\mu(\rho)}. \qquad (31)$$

Here

$$\begin{aligned} \lambda(\rho) \; &= \; \lambda_0 e^{-q\rho(x-a,t)}e^{-m\rho(x+a,t)}, \\ \mu(\rho) \; &= \; \mu_0 e^{-q\rho(x+a,t)}e^{-m\rho(x-a,t)}, \end{aligned} \qquad (32)$$

where the factor $e^{-m\rho(x+a,t)}$ is used to model volume filling: the jump probability to the right $r(\rho)$ is reduced by the the presence of cells at the point $x+a$; m is the volume filling parameter. The factor $e^{-q\rho(x-a,t)}$ describes the adhesion effect, which says that the jump probability to the right $r(\rho)$ is reduced by the presence of cells at the point $x-a$; q is the adhesion coefficient. The master equation for the mean density takes the form

$$\begin{aligned}\frac{\partial \rho}{\partial t} &= \frac{l(\rho(x-a,t))}{{\tau_0}^{\mu(x-a)}}\mathcal{D}_t^{1-\mu(x-a)}\rho\left(x-a,t\right)+\\ &\quad \frac{r(\rho(x+a,t))}{{\tau_0}^{\mu(x-a)}}\mathcal{D}_t^{1-\mu(x+a)}\rho\left(x+a,t\right)-\\ &\quad -\mathcal{D}_t^{1-\mu(x)}\rho\left(x,t\right). \end{aligned}\qquad (33)$$

III. TRANSITION TO ASYMPTOTIC ADVECTION-DIFFUSION TRANSPORT REGIME AND AGGREGATION PHENOMENON

In this section we discuss a transition from a subdiffusive transport regime to an asymptotically normal advection-diffusion transport regime. This transition is governed by the nonlinear fractional equation (25). It involves the exponential factor $e^{-\Phi(x,t)}$, with $\Phi\left(x,t\right)=\int_0^t \alpha\left(\rho\left(x,s\right)\right)ds$, that can be considered as a nonlinear tempering. It generalizes the linear tempering, where the power law waiting time distribution is truncated by an exponential factor $\exp(-\alpha t)$ [35]. We expect that in the limit $\Phi\left(x,t\right)\to\infty$ we recover the classical Fokker-Planck equation, while in the intermediate asymptotic regime, when $\Phi\left(x,t\right)<<1$ we have a subdiffusive transport. If we take the limit $\Phi\left(x,t\right)\to 0$ in (25) for small α, we obtain the standard fractional Fokker-Planck equation [31, 32]

$$\begin{aligned}\frac{\partial \rho}{\partial t} = -\frac{\partial}{\partial x}\left[2D_\mu(x)\beta\frac{\partial U}{\partial x}\mathcal{D}_t^{1-\mu(x)}\rho\right]\\ +\frac{\partial^2}{\partial x^2}\left[D_\mu(x)\mathcal{D}_t^{1-\mu(x)}\rho\right],\end{aligned}\qquad (34)$$

where $D_\mu(x)$ is defined by (26). Nonuniform distribution of the anomalous exponent $\mu(x)$ in (34) leads to the aggregation of particles at the point of minimum of $\mu(x)$ [19]. This phenomenon has been observed in experiments on phagotrophic protists when "cells become immobile in attractive patches, which will then eventually trap all cells" [35]. In this case the potential field $U(x)$ is irrelevant to the distribution of ρ, and anomalous exponent $\mu(x)$ dominates. However, this aggregation around one point is just a transient phenomenon. If we consider the long time limit $t\to\infty$ then we obtain the classical Fokker-Planck equation (see next two subsections). The stationary solution of this equation represents ultimate cells aggregation determined by the potential field $U(x)$ and the spatial distribution of anomalous exponent $\mu(x)$.

A. Aggregation in the linear case

Firstly, we assume that the local escape rate α is independent of time, such that $\Phi\left(x,t\right)=\alpha(x)t$. The Laplace transform of $i\left(x,t\right)$, defined by (19), gives

$$i\left(x,s\right)=\frac{\left(s+\alpha(x)\right)^{1-\mu(x)}}{\tau_0^{\mu(x)}}\rho\left(x,s\right)+\alpha(x)\rho(x,s). \qquad (35)$$

In the limit $s\to 0$ $(t\to\infty)$ the dependence of the first term in (35) on s disappears, so the total escape rate $i\left(x,t\right)$ can be written in the Markovian form

$$i\left(x,t\right)=\lambda_\mu(x)\rho\left(x,t\right),$$

where the effective rate of escape $\lambda_\mu(x)$ is

$$\lambda_\mu(x)=\frac{\alpha(x)}{\left(\tau_0\alpha(x)\right)^{\mu(x)}}+\alpha(x).$$

The essential feature of this rate parameter $\lambda_\mu(x)$ is that it depends on the fractal exponent $\mu(x)$. This escape rate, together with (24), leads to the standard Fokker-Planck equation

$$\begin{aligned}\frac{\partial \rho}{\partial t} &= -\frac{\partial}{\partial x}\left[2\beta\frac{\partial U}{\partial x}D\left(x\right)\rho(x,t)\right]\\ &\quad +\frac{\partial^2}{\partial x^2}\left[D\left(x\right)\rho(x,t)\right],\end{aligned}\qquad (36)$$

where the diffusion coefficient $D\left(x\right)$ depends on $\mu(x)$ and $\alpha(x)$:

$$D\left(x\right)=\frac{a^2\lambda_\mu(x)}{2}=\frac{a^2\left[\left(\tau_0\alpha(x)\right)^{1-\mu(x)}+\tau_0\alpha(x)\right]}{2\tau_0}.$$

When the product $\tau_0\alpha$ is small, the term $\left(\tau_0\alpha\right)^{1-\mu(x)}$ is dominant in the anomalous case $\mu(x)<1$, so $D_\mu\left(x\right)$ can be approximated as

$$D\left(x\right)=\frac{a^2\left(\tau_0\alpha(x)\right)^{1-\mu(x)}}{2\tau_0}.$$

Let us find the stationary solution $\rho_{st}(x)$ to (36) in the interval $[0,L]$. We use the reflective boundary conditions at $x=0$ and $x=L$ which guarantees the conservation of the total population. We introduce the new function

$$p(x)=D\left(x\right)\rho_{st}(x).$$

It follows from (36) that this function obeys the equation

$$\frac{\partial}{\partial x}\left[2\beta\frac{\partial U(x)}{\partial x}p(x)\right]+\frac{\partial^2 p(x)}{\partial x^2}=0 \qquad (37)$$

with the solution in the form of the Boltzmann distribution

$$p(x)=N^{-1}\exp\left[-2\beta U(x)\right].$$

Thus, the steady profile is

$$\rho_{st}(x)=N^{-1}D^{-1}\left(x\right)\exp\left[-2\beta U(x)\right], \qquad (38)$$



FIG. 1: Stationary profiles $\rho_{st}(x)$ for the linear distribution of the potential $U(x) = 5x$ (solid line) and $U(x) = 0$ (dashed line), $\beta = 1$, $\mu_0 = 0.9$, $\alpha = 10^{-4}$, $k = 2.19$.

where N is determined by the normalization condition $N = \int_0^L D^{-1}(x) \exp\left[-2\beta U(x)\right] dx$.

To illustrate how the nonhomogeneous anomalous exponent $\mu(x)$ affects the aggregation pattern we consider the steady profile (38) in the interval $[0, 1]$ for two cases: (a) $U(x) = 0$ and (b) $U(x) = mx$ for which the exponent $\mu(x)$ has the form

$$\mu(x) = \mu_0 \exp\left(-kx\right),$$

where $\mu_0 < 1$. Fig. 1 shows that for the uniform potential $U(x) = 0$, the particles have the tendency to aggregate in the region of the small values of $\mu(x)$ (dashed line). When the potential field forces the particles to move from the right to the left ($U(x) = 5x$), the steady profile is not monotonic (solid line).

B. Aggregation in the nonlinear case

Now we consider the nonlinear case when the escape rate α depends on the density ρ. For living systems, this nonlinear dependence results from a coupling between the local density of cells and the intensity of the response of individual cell to external signals. We assume that in the limit $t \to \infty$ the stationary distribution $\rho_{st}(x)$ exists. Then as $t \to \infty$, the tempering factor $e^{-\Phi(x,t)}$ can be approximated by

$$e^{-\alpha(\rho_{st}(x))t}.$$

The stationary escape rate $i_{st}(x)$ corresponding to (17) can be written in terms of the Laplace transform $\hat{K}(x, s)$ as follows

$$i_{st}(x) = \left[\hat{K}\left(x, \alpha(\rho_{st})\right) + \alpha(\rho_{st})\right] \rho_{st}(x). \qquad (39)$$

This steady rate $i_{st}(x)$ has the Markovian form in which the rate parameter consists of two terms $\hat{K}\left(x, \alpha(\rho_{st})\right)$ and $\alpha(\rho_{st})$. The dependence of the first term on $\alpha(\rho_{st})$ is due to the non-Markovian character of transport process. This effect does not exist in the Markovian case for which $\hat{K}$ is a function of x only.

For the subdiffusive case, the stationary escape rate $i_{st}(x)$ is

$$i_{st}(x) = \left[\frac{\left[\alpha(\rho_{st}(x))\right]^{1-\mu(x)}}{\tau_0^{\mu(x)}} + \alpha(\rho_{st}(x))\right] \rho_{st}(x). \qquad (40)$$

One can see that the first term on the RHS of (40) is dominant for small $\alpha\tau_0$ and $\mu(x) < 1$. For the jump density (21), in the limit $a \to 0$ we obtain the stationary nonlinear Fokker-Planck equation

$$\frac{\partial}{\partial x}\left[2\beta \frac{\partial U}{\partial x} D\left(\rho_{st}\right) \rho_{st}(x)\right] = \frac{\partial^2}{\partial x^2}\left[D\left(\rho_{st}\right) \rho_{st}(x)\right], \qquad (41)$$

where $D\left(\rho_{st}(x)\right)$ is the nonlinear diffusion coefficient defined as

$$D\left(\rho_{st}(x)\right) = \frac{a^2 \left[\alpha(\rho_{st}(x))\right]^{1-\mu(x)}}{2\tau_0^{\mu(x)}}.$$

If we assume a zero flux condition at the boundaries of the interval $[0, L]$, then

$$J = 2\beta \frac{\partial U}{\partial x} D\left(\rho_{st}\right) \rho_{st}(x) - \frac{\partial}{\partial x}\left[D\left(\rho_{st}\right) \rho_{st}(x)\right] = 0, \qquad (42)$$

and the stationary profile $\rho_{st}(x)$ can be found from the nonlinear equation

$$\rho_{st}(x) = N^{-1} D^{-1}\left(\rho_{st}(x)\right) \exp\left[-2\beta U(x)\right]. \qquad (43)$$

The time evolution of the density profile $\rho(x, t)$ for the nonlinear fractional equation (25) can be described as follows. At lower values of $\Phi = \int_0^t \alpha(\rho(x, s)) ds$, the early evolution is the development of a single peak at the point of the minimum of $\mu(x)$. This can be considered as an intermediate anomalous aggregation of particles. However, incorporating the escape rate $\alpha\left(\rho\right)$ and the nonlinear tempering factor $e^{-\Phi}$ provide a regularization of anomalous aggregation. For sufficiently large Φ the density profile $\rho\left(x, t\right)$ must converge to a stationary solution (43) as $t \to \infty$.

IV. CONCLUSIONS.

The aim in this paper was to derive the macroscopic nonlinear subdiffusive fractional equations for the evolution of a mean density of random walkers by incorporating a nonlinear escape rate and nonlinear jump distributions. The main motivation was to take into account the interaction between particles on the mesoscopic level at which the random walker characteristics depend on the mean field density of particles. We showed that the nonlinear escape rate leads to the effective regularization of standard subdiffusive fractional equations. Our modified fractional equations describe the transition from an intermediate subdiffusive regime to an asymptotically normal advection-diffusion transport regime. We showed that this transition is governed by a nonlinear tempering factor that generalize the standard linear tempering. We discussed the aggregation phenomenon and showed the impact of a nonuniform distribution of anomalous exponent on aggregation patterns.

Acknowledgements. The author gratefully acknowledges the support of the EPSRC Grant EP/J019526/1 and the warm hospitality of Department of Mathematical Physics, Ural Federal University. He wishes to thank Steven Falconer, Peter Straka and Andrey Zubarev for interesting discussions.

[1] R. Metzler and J. Klafter, Phys. Rep. 339 1 (2000).

[2] *Anomalous transport: foundations and applications.* Eds. R. Klages, G. Radons, I. M. Sokolov (Wiley-VCH, 2008).

[3] V. Méndez, S. Fedotov and W. Horsthemke, *Reaction-Transport Systems* (Springer, Berlin 2010).

[4] M. J. Saxton, Biophys. J. **81**, 2226 (2001).

[5] I. Golding, E. Cox, Phys. Rev. Lett. **96,** 098102 (2006).

[6] F. Santamaria, S. Wils, E. De Schutter, and G. J. Augustine, Neuron **52**, 635 (2006); S. Fedotov and V. Méndez, Phys. Rev. Lett. **101**, 218102 (2008).

[7] M.B. Miller and B. L. Bassler, Annu. Rev. Microbiol., **55** 165, (2001).

[8] R. A. Weinberg, The Biology of Cancer (Garland Science, New York, 2007).

[9] K. Anguige and C. Schmeiser, J. Math. Biol. **58**, 395 (2009).

[10] S. T. Johnston, M. J. Simpson, and R. E. Baker, Phys. Rev. E **85**, 051922 (2012).

[11] E. Khain, M. Katakowski, S. Hopkins, A. Szalad, X. Zheng, F. Jiang, and M. Chopp, Phys. Rev. E **83**, 031920 (2011).

[12] K. Painter and Th. Hillen, Canadian Applied Mathematics Quarterly, **10** 501 (2002).

[13] V. Méndez, D. Campos, I. Pagonabarraga, and S. Fedotov, J. Theor. Biology, **309** 113 (2012).

[14] S. Fedotov, D. Moss, and D. Campos, Phys. Rev. E **78**, 026107 (2008).

[15] M. O. Vlad and J. Ross, Phys. Rev. E **66**, 061908 (2002).

[16] A. Yadav and W. Horsthemke, Phys. Rev. E **74**, 066118 (2006).

[17] Y. Nec and A. A. Nepomnyashchy, J. Phys. A: Math. Theor. **40** 14687 (2007).

[18] D. Froemberg and I. M. Sokolov, Phys. Rev. Lett. **100**, 108304 (2008).

[19] S. Fedotov, Phys. Rev. E **81**, 011117 (2010).

[20] E. Abad, S. B. Yuste, and K. Lindenberg, Phys. Rev. E **81**, 031115 (2010).

[21] S. Fedotov, A. Iomin, Phys. Rev. Lett.*,* **98** 118101 (2007); S. Fedotov, Phys. Rev. E **81**, 011117 (2010).

[22] F. Sagues, V. P. Shkilev, I. M. Sokolov, Phys. Rev. E **77** (2008), 032102.

[23] V. P. Shkilev. J. Exp. Theor. Physics, **112**, 711 (2011).

[24] V. A. Volpert, Y. Nec, A. A. Nepomnyashchy. Phil. Trans. R. Soc. A **371**, 20120179 (2013).

[25] S. Fedotov, Phys. Rev. E **83**, 021110 (2011).

[26] S. Fedotov and S. Falconer, Phys. Rev. E **85**, 031132 (2012).

[27] D. R. Cox and H. D. Miller, *The Theory of Stochastic Processes* (Methuen, London, 1965)

[28] E. Scalas, R. Gorenflo, F. Mainardi, and M. Raberto, Fractals **11**, 281 (2003).

[29] A. V. Chechkin, R. Gorenflo and I. M. Sokolov, J. Phys. A: Math. Gen **38**, L679 (2005).

[30] S. Fedotov and S. Falconer, arXiv:1210.8020 [cond-mat.stat-mech] (2012).

[31] E. Barkai, R. Metzler, and J. Klafter, Phys. Rev. E **61**, 132 (2000).

[32] B. I. Henry, T. A. M. Langlands, and P. Straka, Phys. Rev. Lett. **105,** 170602 (2010).

[33] H. G. Othmer and A. Stevens, SIAM J. Appl. Math. **57**, 1044 (1997).

[34] T. A. M. Langlands and B. I. Henry, Phys. Rev. E **81**, 051102 (2010).

[35] M. M. Meerschaert, Geophys. Res. Lett. 35, L17403 (2008); A. I. Saichev, and S. G. Utkin , J. Exp. Theory Phys., **99**, 443 (2004); A. Stanislavsky, K. Weron, and A. Weron, Phys. Rev. E **78**, 051106 (2008); J. Gajda and M. Magdziarz, Phys. Rev. E **82**, 011117 (2010).

[36] T. Fenchel and N. Blackburn, Protist **160**, 325 (1999).